arXiv:1604.01852v1 [math.CA] 7 Apr 2016

NEW INEQUALITIES OF STEFFENSEN'S TYPE FOR s–CONVEX FUNCTIONS

MOHAMMAD W. ALOMARI

Abstract. In this work, new inequalities connected with the Steffensen's integral inequality for s-convex functions are proved.

1. Introduction

In order to study certain inequalities between mean values, J.F. Steffensen [11] has proved the following inequality (see also [9], p. 311):

Theorem 1. *Let f and g be two integrable functions defined on (a,b), f is decreasing and for each $t\in(a,b)$, $0\leq g(t)\leq 1$. Then, the following inequality*

$$\int_{b-\lambda}^{b} f(t)\,dt \leq \int_{a}^{b} f(t)\,g(t)\,dt \leq \int_{a}^{a+\lambda} f(t)\,dt \tag{1.1}$$

holds, where, $\lambda=\int_a^b g(t)\,dt$.

Some minor generalization of Steffensen's inequality (1.1) was considered by T. Hayashi [5], using the substituting $g(t)/A$ for $g(t)$, where A is positive constant. For other result involving Steffensen type inequality, see [3], [5] and [8]–[11].

In the recent work [1], Alomari et. al. proved the following result:

Theorem 2. *Let $f,g:[a,b]\to\mathbb{R}$ be integrable such that $0\leq g(t)\leq 1$, for all $t\in[a,b]$ such that $\int_a^b g(t)\,df(t)$ exists. If f is absolutely continuous on $[a,b]$ with $f'\in L_p[a,b]$, $1\leq p\leq\infty$, then we have*

$$\left|\int_a^{a+\lambda} f(t)\,dt-\int_a^b f(t)\,g(t)\,dt\right| \tag{1.2}$$
$$\leq\begin{cases} \frac{1}{2}\left[\lambda^2+(b-a-\lambda)^2\right]\|f'\|_{\infty,[a,b]}, & if \quad f'\in L_\infty[a,b];\\ \frac{\|f'\|_{p,[a,b]}}{(q+1)^{1/q}}\left[\lambda^{(q+1)/q}+(b-a-\lambda)^{(q+1)/q}\right], & if \quad f'\in L_p[a,b],p>1;\\ \left[\int_{a+\lambda}^b g(t)\,dt\right]\|f'\|_1, & if \quad f'\in L_1[a,b], \end{cases}$$

2000 *Mathematics Subject Classification.* 26D10, 26D15.
Key words and phrases. Steffensen inequality, Hayashi's inequality, s–Convex function.

and

$$\left|\int_a^b f(t)\,g(t)\,dt-\int_{b-\lambda}^b f(t)\,dt\right| \tag{1.3}$$
$$\leq\begin{cases}\frac{1}{2}\left[\lambda^2+(b-a-\lambda)^2\right]\|f'\|_{\infty,[a,b]}, & if \quad f'\in L_\infty[a,b];\\ \frac{\|f'\|_{p,[a,b]}}{(q+1)^{1/q}}\left[\lambda^{(q+1)/q}+(b-a-\lambda)^{(q+1)/q}\right], & if \quad f'\in L_p[a,b],p>1;\\ \left[\int_a^{b-\lambda}g(t)\,dt\right]\|f'\|_1, & if \quad f'\in L_1[a,b].\end{cases}$$

A function $f:\mathbb{R}^+\to\mathbb{R}$, where $\mathbb{R}^+=[0,\infty)$, is said to be s-convex in the second sense if

$$f(\alpha x+\beta y)\leq\alpha^s f(x)+\beta^s f(y)$$

for all $x,y\in[0,\infty)$, $\alpha,\beta\geq 0$ with $\alpha+\beta=1$ and for some fixed $s\in(0,1]$. This class of s-convex functions is usually denoted by K_s^2, (see [6]). It can be easily seen that for $s=1$, s-convexity reduces to the ordinary convexity of functions defined on $[0,\infty)$.

In [4], Dragomir and Fitzpatrick proved a variant of Hadamard's inequality which holds for s–convex functions in the second sense:

$$2^{s-1}f\left(\frac{a+b}{2}\right)\leq\frac{1}{b-a}\int_a^b f(x)\,dx\leq\frac{f(a)+f(b)}{s+1}. \tag{1.4}$$

The constant $k=\frac{1}{s+1}$ is the best possible in the second inequality in (1.4). For another inequalities of Hermite–Hadamard type see [2] and [7].

The aim of this paper is to establish new inequalities of Steffensen's type for s-convex functions in the second sense.

2. The results

Lemma 1. *Let $f,g:[a,b]\subset\mathbb{R}^+\to\mathbb{R}$ be integrable such that $0\leq g(t)\leq 1$, for all $t\in[a,b]$ and $\int_a^b g(t)\,f'(t)\,dx$ exists. Then we have the following representation*

$$\int_a^{a+\lambda}f(t)\,dt-\int_a^b f(t)\,g(t)\,dt \tag{2.1}$$
$$=-\int_a^{a+\lambda}\left(\int_a^x(1-g(t))\,dt\right)f'(x)\,dx-\int_{a+\lambda}^b\left(\int_x^b g(t)\,dt\right)f'(x)\,dx,$$

and

$$\int_a^b f(t)\,g(t)\,dt-\int_{b-\lambda}^b f(t)\,dt \tag{2.2}$$
$$=-\int_a^{b-\lambda}\left(\int_a^x g(t)dt\right)f'(x)\,dx-\int_{b-\lambda}^b\left(\int_x^b(1-g(x))\,dt\right)f'(x)\,dx,$$

where $\lambda:=\int_a^b g(t)\,dt$.

Proof. Integrating by parts

$$
\begin{aligned}
&-\int_a^{a+\lambda}\left(\int_a^x (1-g(t))\,dt\right) f'(x)\,dx-\int_{a+\lambda}^b\left(\int_x^b g(t)\,dt\right) f'(x)\,dx \\
&=-\left(\int_a^{a+\lambda}(1-g(t))\,dt\right) f(a+\lambda)+\int_a^{a+\lambda} f(x)\,d\left(\int_a^x (1-g(t))\,dt\right) \\
&\quad+\left(\int_{a+\lambda}^b g(t)\,dt\right) f(a+\lambda)+\int_{a+\lambda}^b f(x)\,d\left(\int_x^b g(t)\,dt\right) \\
&=-\left(\int_a^{a+\lambda}(1-g(t))\,dt\right) f(a+\lambda)+\int_a^{a+\lambda} f(x)\,(1-g(x))\,dx \\
&\quad+\left(\int_{a+\lambda}^b g(t)\,dt\right) f(a+\lambda)-\int_{a+\lambda}^b f(x)\,g(x)\,dx \\
&=-\lambda f(a+\lambda)+f(a+\lambda)\int_a^{a+\lambda} g(t)\,dt+\int_a^{a+\lambda} f(x)\,dx \\
&\quad-\int_a^{a+\lambda} f(x)\,g(x)\,dx+f(a+\lambda)\int_{a+\lambda}^b g(t)\,dt-\int_{a+\lambda}^b f(x)\,g(x)\,dx \\
&=-\lambda f(a+\lambda)+f(a+\lambda)\int_a^{a+\lambda} g(t)\,dt+f(a+\lambda)\int_{a+\lambda}^b g(t)\,dt \\
&\quad+\int_a^{a+\lambda} f(x)\,dx-\int_a^b f(x)\,g(x)\,dx \\
&=\int_a^{a+\lambda} f(x)\,dx-\int_a^b f(x)\,g(x)\,dx,
\end{aligned}
$$

which gives the desired representation (2.1). The identity (2.2) can be also proved in a similar way, we shall omit the details. □

2.1. **Inequalities involving s-convexity.** In the following, inequalities for absolutely continuous functions whose first derivatives are s-convex (s-concave) are given:

Theorem 3. *Let $f,g:[a,b]\subset\mathbb{R}^+\to\mathbb{R}$ be integrable such that $0\le g(t)\le 1$, for all $t\in[a,b]$ such that $\int_a^b g(t)\,f'(t)\,dx$ exists. If f is absolutely continuous on $[a,b]$ such that $|f'|$ is s-convex on $[a,b]$, then*

$$
\begin{aligned}
(2.3)\quad &\left|\int_a^{a+\lambda} f(t)\,dt-\int_a^b f(t)\,g(t)\,dt\right| \\
&\le\frac{1}{(s+1)(s+2)}\left[\lambda^2|f'(a)|+(b-a-\lambda)^2|f'(b)|\right] \\
&\qquad+\frac{1}{s+2}\left[\lambda^2+(b-a-\lambda)^2\right]|f'(a+\lambda)|,
\end{aligned}
$$

and

$$\left|\int_a^b f(t)\,g(t)\,dt-\int_{b-\lambda}^{b} f(t)\,dt\right| \le \frac{1}{(s+1)(s+2)}\left[\lambda^2\left|f'(b)\right|+(b-a-\lambda)^2\left|f'(a)\right|\right] + \frac{1}{s+2}\left[\lambda^2+(b-a-\lambda)^2\right]\left|f'(b-\lambda)\right|, \tag{2.4}$$

where, $\lambda := \int_a^b g(t)\,dt$.

Proof. Utilizing the triangle inequality on (2.1), and since $|f'|$ is s-convex, we have

$$\begin{aligned}
&\left|\int_a^{a+\lambda} f(t)\,dt-\int_a^b f(t)\,g(t)\,dt\right| \\
&\le \left|\int_a^{a+\lambda}\left(\int_a^x (1-g(t))\,dt\right)f'(x)\,dx\right|+\left|\int_{a+\lambda}^{b}\left(\int_x^b g(t)\,dt\right)f'(x)\,dx\right| \\
&\le \int_a^{a+\lambda}\left|\int_a^x (1-g(t))\,dt\right|\left|f'(x)\right|dx+\int_{a+\lambda}^{b}\left|\int_x^b g(t)\,dt\right|\left|f'(x)\right|dx \\
&\le \int_a^{a+\lambda}\left|\int_a^x (1-g(t))\,dt\right|\left[\frac{(x-a)^s}{\lambda^s}\left|f'(a+\lambda)\right|+\frac{(a+\lambda-x)^s}{\lambda^s}\left|f'(a)\right|\right]dx \\
&\quad+\int_{a+\lambda}^{b}\left|\int_x^b g(t)\,dt\right|\left[\frac{(x-a-\lambda)^s}{(b-a-\lambda)^s}\left|f'(b)\right|+\frac{(b-x)^s}{(b-a-\lambda)^s}\left|f'(a+\lambda)\right|\right]dx \\
&\le \frac{\left|f'(a+\lambda)\right|}{\lambda^s}\int_a^{a+\lambda}\left(\int_a^x\left|1-g(t)\right|dt\right)(x-a)^s\,dx \\
&\quad+\frac{\left|f'(a)\right|}{\lambda^s}\int_a^{a+\lambda}\left(\int_a^x\left|1-g(t)\right|dt\right)(a+\lambda-x)^s\,dx \\
&\quad+\frac{\left|f'(b)\right|}{(b-a-\lambda)^s}\int_{a+\lambda}^{b}\left(\int_x^b\left|g(t)\right|dt\right)(x-a-\lambda)^s\,dx \\
&\quad+\frac{\left|f'(a+\lambda)\right|}{(b-a-\lambda)^s}\int_{a+\lambda}^{b}\left(\int_x^b\left|g(t)\right|dt\right)(b-x)^s\,dx \\
&\le \frac{\left|f'(a+\lambda)\right|}{\lambda^s}\int_a^{a+\lambda}(x-a)^{s+1}\,dx+\frac{\left|f'(a)\right|}{\lambda^s}\int_a^{a+\lambda}(x-a)(a+\lambda-x)^s\,dx \\
&\quad+\frac{\left|f'(b)\right|}{(b-a-\lambda)^s}\int_{a+\lambda}^{b}(b-x)(x-a-\lambda)^s\,dx+\frac{\left|f'(a+\lambda)\right|}{(b-a-\lambda)^s}\int_{a+\lambda}^{b}(b-x)^{s+1}\,dx \\
&= \frac{1}{(s+1)(s+2)}\left[\lambda^2\left|f'(a)\right|+(b-a-\lambda)^2\left|f'(b)\right|\right] \\
&\quad+\frac{1}{s+2}\left[\lambda^2+(b-a-\lambda)^2\right]\left|f'(a+\lambda)\right|
\end{aligned}$$

which proves the first inequality in (2.3). In similar way and using (2.2) we may deduce the desired inequality (2.4), and we shall omit the details. □

Corollary 1. *In (2.3) if one chooses $s = 1$ then*

$$\begin{aligned}(2.5)\quad &\left|\int_a^{a+\lambda} f(t)\,dt - \int_a^b f(t)\,g(t)\,dt\right| \\ &\leq \frac{1}{6}\lambda^2\left|f'(a)\right| + \frac{1}{3}\left[\lambda^2 + (b-a-\lambda)^2\right]\left|f'(a+\lambda)\right| + \frac{1}{6}(b-a-\lambda)^2\left|f'(b)\right|\end{aligned}$$

also, in (2.4) if $s = 1$, then

$$\begin{aligned}(2.6)\quad &\left|\int_a^b f(t)\,g(t)\,dt - \int_{b-\lambda}^b f(t)\,dt\right| \\ &\leq \frac{1}{6}\lambda^2\left|f'(b)\right| + \frac{1}{3}\left[\lambda^2 + (b-a-\lambda)^2\right]\left|f'(b-\lambda)\right| + \frac{1}{6}(b-a-\lambda)^2\left|f'(a)\right|\end{aligned}$$

Remark 1. *In the inequalities (2.3) and (2.4), choose $\lambda = 0$, then we have*

$$\begin{aligned}(2.7)\quad &\left|\int_a^b f(t)\,g(t)\,dt\right| \\ &\leq \frac{(b-a)^2}{(s+1)(s+2)}\min\left\{(s+1)\left|f'(a)\right| + \left|f'(b)\right|, \left|f'(a)\right| + (s+1)\left|f'(b)\right|\right\}.\end{aligned}$$

Another approach leads to the following result:

Theorem 4. *Let $f, g : [a,b] \subset \mathbb{R}^+ \to \mathbb{R}$ be integrable such that $0 \leq g(t) \leq 1$, for all $t \in [a,b]$ such that $\int_a^b g(t)\,f'(t)\,dx$ exists. If f is absolutely continuous on $[a,b]$ with $|f'|$ is s-convex on $[a,b]$, for some fixed $s \in (0,1]$ then we have*

$$\begin{aligned}(2.8)\quad &\left|\int_a^{a+\lambda} f(t)\,dt - \int_a^b f(t)\,g(t)\,dt\right| \\ &\leq \frac{1}{s+1}\left[\int_{a+\lambda}^b g(t)\,dt\right]\cdot\left[\lambda\left|f'(a)\right| + (b-a)\left|f'(a+\lambda)\right| + (b-a-\lambda)\left|f'(b)\right|\right] \\ &\leq \frac{(b-a-\lambda)}{s+1}\left[\lambda\left|f'(a)\right| + (b-a)\left|f'(a+\lambda)\right| + (b-a-\lambda)\left|f'(b)\right|\right],\end{aligned}$$

and

$$\begin{aligned}(2.9)\quad &\left|\int_a^b f(t)\,g(t)\,dt - \int_{b-\lambda}^b f(t)\,dt\right| \\ &\leq \frac{1}{s+1}\left[\int_{b-\lambda}^b g(t)\,dt\right]\cdot\left[(b-a-\lambda)\left|f'(a)\right| + (b-a)\left|f'(b-\lambda)\right| + \lambda\left|f'(b)\right|\right] \\ &\leq \frac{\lambda}{s+1}\left[(b-a-\lambda)\left|f'(a)\right| + (b-a)\left|f'(b-\lambda)\right| + \lambda\left|f'(b)\right|\right],\end{aligned}$$

where $\lambda := \int_a^b g(t)\,dt$.

Proof. From Lemma 1, we may write

$$\left|\int_a^{a+\lambda} f(t)\,dt - \int_a^b f(t)\,g(t)\,dt\right|$$
$$\leq \sup_{x\in[a,a+\lambda]}\left[\int_a^x (1-g(t))\,dt\right]\cdot\int_a^{a+\lambda}|f'(x)|\,dx + \sup_{x\in[a+\lambda,b]}\left[\int_x^b g(t)\,dt\right]\cdot\int_{a+\lambda}^b |f'(x)|\,dx.$$

Since $|f'|$ is s–convex on $[a,b]$, then by (1.4) we have

$$\int_a^{a+\lambda}|f'(x)|\,dx \leq \lambda\cdot\frac{|f'(a)|+|f'(a+\lambda)|}{s+1},$$

and

$$\int_{a+\lambda}^b |f'(x)|\,dx \leq (b-a-\lambda)\cdot\frac{|f'(a+\lambda)|+|f'(b)|}{s+1}.$$

Therefore, we have

$$\begin{aligned}
&\left|\int_a^{a+\lambda} f(t)\,dt - \int_a^b f(t)\,g(t)\,dt\right| \\
&\leq \lambda\cdot\frac{|f'(a)|+|f'(a+\lambda)|}{s+1}\cdot\int_a^{a+\lambda}(1-g(t))\,dt \\
&\quad + (b-a-\lambda)\cdot\frac{|f'(a+\lambda)|+|f'(b)|}{s+1}\cdot\int_{a+\lambda}^b g(t)\,dt \\
&\leq \max\left\{\int_a^{a+\lambda}(1-g(t))\,dt, \int_{a+\lambda}^b g(t)\,dt\right\}\cdot\left[\lambda\frac{|f'(a)|+|f'(a+\lambda)|}{s+1}\right. \\
&\quad \left. + (b-a-\lambda)\,\frac{|f'(a+\lambda)|+|f'(b)|}{s+1}\right] \\
&= \frac{1}{s+1}\left[\int_{a+\lambda}^b g(t)\,dt\right]\cdot\left[\lambda|f'(a)| + (b-a)|f'(a+\lambda)| + (b-a-\lambda)|f'(b)|\right],
\end{aligned}$$

which proves the first inequality in (2.8). The second inequality in (2.8) follows directly, since $0\leq g(t)\leq 1$ for all $t\in[a,b]$, then

$$0\leq\int_{a+\lambda}^b g(t)\,dt\leq(b-a-\lambda).$$

The inequalities in (2.9) may be proved in the same way using the identity (2.2), we shall omit the details. □

2.2. **Inequalities involving s-concavity.**

Theorem 5. *Let $f,g:[a,b]\subset\mathbb{R}^+\to\mathbb{R}$ be integrable such that $0\leq g(t)\leq 1$, for all $t\in[a,b]$ such that $\int_a^b g(t)\,f'(t)\,dx$ exists. If f is absolutely continuous on $[a,b]$*

with $|f'|$ is s-concave on $[a,b]$, for some fixed $s\in(0,1]$ then we have

$$\begin{aligned}
&\left|\int_a^{a+\lambda} f(t)\,dt-\int_a^b f(t)\,g(t)\,dt\right| \\
&\leq 2^{s-1}\left[\int_{a+\lambda}^b g(t)\,dt\right]\cdot\left[\lambda\left|f'\left(a+\frac{\lambda}{2}\right)\right|+(b-a-\lambda)\left|f'\left(\frac{a+b+\lambda}{2}\right)\right|\right]. \\
&\leq 2^{s-1}(b-a-\lambda)\cdot\left[\lambda\left|f'\left(a+\frac{\lambda}{2}\right)\right|+(b-a-\lambda)\left|f'\left(\frac{a+b+\lambda}{2}\right)\right|\right]
\end{aligned} \tag{2.10}$$

and

$$\begin{aligned}
&\left|\int_a^b f(t)\,g(t)\,dt-\int_{b-\lambda}^b f(t)\,dt\right| \\
&\leq 2^{s-1}\left[\int_{b-\lambda}^b g(t)\,dt\right]\cdot\left[(b-a-\lambda)\left|f'\left(\frac{a+b-\lambda}{2}\right)\right|+\lambda\left|f'\left(b-\frac{\lambda}{2}\right)\right|\right]. \\
&\leq \lambda 2^{s-1}\cdot\left[(b-a-\lambda)\left|f'\left(\frac{a+b-\lambda}{2}\right)\right|+\lambda\left|f'\left(b-\frac{\lambda}{2}\right)\right|\right],
\end{aligned} \tag{2.11}$$

where $\lambda:=\int_a^b g(t)\,dt$.

Proof. Utilizing the triangle inequality on (2.1), and since $|f'|$ is s–concave on $[a,b]$ then by (1.4) we may state

$$\begin{aligned}
&\left|\int_a^{a+\lambda} f(t)\,dt-\int_a^b f(t)\,g(t)\,dt\right| \\
&\leq \sup_{x\in[a,a+\lambda]}\left[\int_a^x(1-g(t))\,dt\right]\cdot\int_a^{a+\lambda}|f'(x)|\,dx+\sup_{x\in[a+\lambda,b]}\left[\int_x^b g(t)\,dt\right]\cdot\int_{a+\lambda}^b|f'(x)|\,dx \\
&=2^{s-1}\lambda\left|f'\left(a+\frac{\lambda}{2}\right)\right|\cdot\int_a^{a+\lambda}(1-g(t))\,dt+2^{s-1}(b-a-\lambda)\left|f'\left(\frac{a+b+\lambda}{2}\right)\right|\cdot\int_{a+\lambda}^b g(t)\,dt \\
&=2^{s-1}\left[\int_{a+\lambda}^b g(t)\,dt\right]\cdot\left[\lambda\left|f'\left(a+\frac{\lambda}{2}\right)\right|+(b-a-\lambda)\left|f'\left(\frac{a+b+\lambda}{2}\right)\right|\right].
\end{aligned}$$

which proves the first inequality in (2.10). The second inequality in (2.10) follows directly, since $0\leq g(t)\leq 1$ for all $t\in[a,b]$, then

$$0\leq\int_{a+\lambda}^b g(t)\,dt\leq(b-a-\lambda).$$

The inequalities in (2.11) may be proved in the same way using the identity (2.2), we shall omit the details. □

Another result is incorporated in the following theorem:

Theorem 6. *Let $f,g:[a,b]\subset\mathbb{R}^+\to\mathbb{R}$ be integrable such that $0\leq g(t)\leq 1$, for all $t\in[a,b]$ such that $\int_a^b g(t)\,f'(t)\,dx$ exists. If f is absolutely continuous on $[a,b]$*

with $|f'|^q$ is s-concave on $[a,b]$, for some fixed $s \in (0,1]$ and $q>1$, then we have

$$\left|\int_a^{a+\lambda} f(t)\,dt - \int_a^b f(t)\,g(t)\,dt\right| \le 2^{(s-1)/q}\left[\lambda^2\left|f'\left(a+\frac{\lambda}{2}\right)\right| + (b-a-\lambda)^2\left|f'\left(\frac{a+b+\lambda}{2}\right)\right|\right], \tag{2.12}$$

and

$$\left|\int_a^b f(t)\,g(t)\,dt - \int_{b-\lambda}^b f(t)\,dt\right| \le 2^{(s-1)/q}\left[(b-a-\lambda)^2\left|f'\left(b-\frac{\lambda}{2}\right)\right| + \lambda^2\left|f'\left(\frac{a+b-\lambda}{2}\right)\right|\right], \tag{2.13}$$

where $\lambda := \int_a^b g(t)\,dt$.

Proof. From Lemma 1 and using the Hölder inequality for $q>1$, and $p=\frac{q}{q-1}$, we obtain

$$\begin{aligned}
&\left|\int_a^{a+\lambda} f(t)\,dt - \int_a^b f(t)\,g(t)\,dt\right| \\
&\le \int_a^{a+\lambda}\left|\int_a^x (1-g(t))\,dt\right||f'(x)|\,dx + \int_{a+\lambda}^b\left|\int_x^b g(t)\,dt\right||f'(x)|\,dx \\
&\le \left(\int_a^{a+\lambda}\left|\int_a^x (1-g(t))\,dt\right|^p dx\right)^{1/p}\left(\int_a^{a+\lambda}|f'(x)|^q\,dx\right)^{1/q} \\
&\quad + \left(\int_{a+\lambda}^b\left|\int_x^b g(t)\,dt\right|^p dx\right)^{1/p}\left(\int_{a+\lambda}^b |f'(x)|^q\,dx\right)^{1/q} := M,
\end{aligned} \tag{2.14}$$

where p is the conjugate of q.

By the inequality (1.4), we have

$$\int_a^{a+\lambda}|f'(x)|^q\,dx \le 2^{s-1}\lambda\left|f'\left(a+\frac{1}{2}\lambda\right)\right|^q,$$

and

$$\int_{a+\lambda}^b |f'(x)|^q\,dx \le 2^{s-1}(b-a-\lambda)\left|f'\left(\frac{a+b+\lambda}{2}\right)\right|^q,$$

which gives by (2.14)

$$\begin{aligned}
M &\leq 2^{(s-1)/q}\lambda^{1/q}\left|f'\left(a+\frac{\lambda}{2}\right)\right|\left(\int_a^{a+\lambda}(x-a)^p\,dx\right)^{1/p} \\
&\quad + 2^{(s-1)/q}\left(b-a-\lambda\right)^{1/q}\left|f'\left(\frac{a+b+\lambda}{2}\right)\right|\left(\int_{a+\lambda}^{b}(b-x)^p\,dx\right)^{1/p} \\
&= 2^{(s-1)/q}\left[\lambda^{1+\frac{1}{p}+\frac{1}{q}}\left|f'\left(a+\frac{\lambda}{2}\right)\right|+\left(b-a-\lambda\right)^{1+\frac{1}{p}+\frac{1}{q}}\left|f'\left(\frac{a+b+\lambda}{2}\right)\right|\right] \\
&= 2^{(s-1)/q}\left[\lambda^{2}\left|f'\left(a+\frac{\lambda}{2}\right)\right|+\left(b-a-\lambda\right)^{2}\left|f'\left(\frac{a+b+\lambda}{2}\right)\right|\right],
\end{aligned}$$

giving the second inequality in (2.12).

The inequalities in (2.13) may be proved in the same way using the identity (2.2), we shall omit the details. □

Remark 2. *The interested reader may obtain several inequalities for log-convex, quasi-convex, r-convex and h-convex functions by replacing the condition on $|f'|$.*

References

[1] M.W. Alomari, S. Hussian and Z. Liu, Some Steffensen's type inequalities, submitted.

[2] M.W. Alomari, M. Darus and U.S. Kirmaci, Some inequalities of Hermite–Hadamard type for s–convex functions, *Acta Mathematica Scientia*, 31B(4):1643–1652, (2011).

[3] A. Apéry, Une inégalité surles fonctions de variable réelle, Attj del Quarto Congresso dell'Unione Matematica Itliana 1951, 2, 3–4 (1953).

[4] S.S. Dragomir and S. Fitzpatrick, The Hadamard's inequality for s-convex functions in the second sense, *Demonstratio Math.*, 32 (4) (1999), 687-696.

[5] T. Hayashi, On curves with monotonous curvature, Tôhoku Math. J., 15, 236–239 (1919).

[6] H. Hudzik and L. Maligranda, Some remarks on s-convex functions, *Aequationes Math.*, 48 (1994), 100-111.

[7] U.S. Kirmaci et al., Hadamard-type inequalities for s-convex functions, *Appl. Math. Comp.*, 193 (2007), 26–35.

[8] B. Meidell, Note sur quelques inégalités et formules d'approximation, *Skand. Aktuarietids.*, 10, 180–198, (1918).

[9] D.S. Mitrinović, J.E. Pečarić and A.M. Fink, Classical and new inequalities in analysis, Kluwer, Dordrecht (1993).

[10] A. Ostrowski, Aufgabensammlung zur infinitesimalrechnung, Basel–Stutgar, 1, (1964).

[11] J.F. Steffensen, On certain inequalities between mean values and their application to actuarial problems, *Skand. Aktuarietids.*, 82–97, (1918).

Department of Mathematics, Faculty of Science, Jerash University, 26150 Jerash, Jordan

E-mail address: mwomath@gmail.com